

8-1-2006

# The Harmonic Oscillator, The Ladder Operator Solutions

Carl W. David

*University of Connecticut*, [Carl.David@uconn.edu](mailto:Carl.David@uconn.edu)

Follow this and additional works at: [https://opencommons.uconn.edu/chem\\_educ](https://opencommons.uconn.edu/chem_educ)

 Part of the [Chemistry Commons](#)

---

## Recommended Citation

David, Carl W., "The Harmonic Oscillator, The Ladder Operator Solutions" (2006). *Chemistry Education Materials*. 19.  
[https://opencommons.uconn.edu/chem\\_educ/19](https://opencommons.uconn.edu/chem_educ/19)

# The Harmonic Oscillator, The Ladder Operator Solutions

C. W. David

*Department of Chemistry  
University of Connecticut  
Storrs, Connecticut 06269-3060*

(Dated: August 1, 2006)

## I. SYNOPSIS

The solution to the quantum mechanical harmonic oscillator using ladder operators is a classic, whose ideas permeate other problem's treatments.

## II. HAMILTONIANS AND COMMUTATORS

The Hamiltonian for the Harmonic Oscillator is

$$\frac{p^2}{2\mu} + \frac{k}{2}x^2$$

where  $p$  is the momentum operator and  $x$  is the position operator. We know that the Schrödinger prescription is

$$p \rightarrow -i\hbar \frac{\partial}{\partial x}$$

while  $x \rightarrow x$ , as usual. This means, as we well know, that the commutator of  $x$  and  $p$  is non-zero.

$$[p, x]f = (px - xp)f = -i\hbar \frac{\partial}{\partial x}xf - x(-i\hbar \frac{\partial}{\partial x}f)$$

which leads to

$$[p, x]f = (px - xp)f = -f i\hbar \frac{\partial x}{\partial x} - x(i\hbar \frac{\partial}{\partial x})f x(-i\hbar \frac{\partial}{\partial x}f)$$

which means, of course,

$$[p, x] = px - xp = -i\hbar$$

For the Harmonic Oscillator, we form the two operators

$$a^+ = p + i\mu\omega x$$

and

$$a^- = p - i\mu\omega x$$

which differ solely by that intervening sign (Remember that  $\omega = \sqrt{\frac{k}{\mu}}$ ). The entire derivation now hinges on the properties of these two operators. We start with the elementary question, what is the commutator of  $a^+$  and  $a^-$ ?

$$[a^+, a^-] = a^+a^- - a^-a^+$$

by definition; these terms expand to become

$$[a^+, a^-] = (p + i\mu\omega x)(p - i\mu\omega x) - (p - i\mu\omega x)(p + i\mu\omega x)$$

which is

$$[a^+, a^-] = p^2 - p i\mu\omega x + i\mu\omega x p + \mu^2\omega^2 x^2 - (p^2 + p i\mu\omega x - i\mu\omega x p + \mu^2\omega^2 x^2)$$

$$[a^+, a^-] = -i\mu\omega(px - xp + px - xp) = -2i\mu\omega(px - xp) = -2i\mu\omega(-i\hbar) = -2\mu\omega\hbar$$

Now the product  $a^+a^- + a^-a^+$  is

$$2(p^2 + \mu^2\omega^2 x^2)$$

so

$$\frac{a^+a^- + a^-a^+}{4} = \frac{p^2 + \mu^2\omega^2 x^2}{2}$$

i.e.,

$$\frac{a^+a^- + a^-a^+}{4\mu} = \frac{p^2 + \mu^2\frac{k}{\mu}x^2}{2\mu} = H$$

We next need the commutator of these "a" operators and the energy operator. We have

$$[a^+, H] = a^+ H - H a^+ = a^+ \frac{a^+ a^- + a^- a^+}{4\mu} - \frac{a^+ a^- + a^- a^+}{4\mu} a^+$$

which is

$$[a^+, H] = \frac{1}{4\mu} (a^+(a^+ a^- + a^- a^+) - (a^+ a^- + a^- a^+) a^+)$$

$$[a^+, H] = \frac{1}{4\mu} (a^+ a^+ a^- + a^+ a^- a^+ - a^+ a^- a^+ - a^- a^+ a^+)$$

$$[a^+, H] = \frac{1}{4\mu} (a^+(a^+ a^- - a^- a^+) + (a^+ a^- - a^- a^+) a^+)$$

$$[a^+, H] = \frac{-1}{4\mu} (a^+(2\mu\hbar\omega) + (2\mu\hbar\omega)a^+) \quad (2.1)$$

$$[a^+, H] = -\hbar\omega a^+$$

This is a crucial result, since it contains the seed of what a ladder operator does.

### III. COMMUTING THE LADDER OPERATOR WITH THE HAMILTONIAN

Consider if we had in hand an eigenfunction  $\psi$  of  $H$ . That would mean that

$$H\psi = E\psi$$

Now operate from the left with  $a^+$  and let's see what happens. We have

$$a^+(H\psi) = (a^+ H)\psi = E a^+ \psi$$

(operators ignore constants like  $E$ ) which is, employing the commutator  $a^+ H - H a^+ = -\hbar\omega a^+$  (Equation 2.1) we have

$$(-\hbar\omega a^+ + H a^+) \psi = E (a^+ \psi)$$

$$(-\hbar\omega a^+) \psi + (H a^+) \psi = E (a^+ \psi)$$

$$H (a^+ \psi) = E (a^+ \psi) + \hbar\omega (a^+ \psi)$$

$$H (a^+ \psi) = (E + \hbar\omega) (a^+ \psi)$$

What does this last statement say? It says that  $a^+ \psi$  is *also* an eigenfunction of  $H$  if  $\psi$  was. The only thing is that the energy is higher than the energy associated with  $\psi$  (by  $\hbar\omega$ ).

What we have shown is that  $a^+$  is an operator, a ladder operator, which takes an existing eigenfunction, and transforms it into the next highest eigenfunction, separated in energy by  $\hbar\omega$ . Wonderful.

Now we need to see what happens with  $a^-$ . We have

$$[a^-, H] = a^- H + H a^- = a^- \left( \frac{a^- a^+ + a^+ a^-}{4\mu} \right) - \left( \frac{a^- a^+ + a^+ a^-}{4\mu} \right) a^-$$

which is

$$[a^-, H] = \frac{1}{4\mu} (a^-(a^- a^+ + a^+ a^-) - (a^- a^+ + a^+ a^-) a^-)$$

$$[a^-, H] = \frac{1}{4\mu} (a^- a^- a^+ + a^- a^+ a^- - a^- a^+ a^- - a^+ a^- a^-)$$

$$[a^-, H] = \frac{1}{4\mu} (a^-(a^- a^+ - a^+ a^-) - (a^- a^+ - a^+ a^-) a^-)$$

$$[a^-, H] = \frac{1}{4\mu} (a^-(2\mu\hbar\omega) + (2\mu\hbar\omega)a^-)$$

$$[a^-, H] = \hbar\omega a^-$$

Again, given

$$H\psi = E\psi$$

Now operate from the left with  $a^-$  and let's see what happens. We have

$$(a^-) H \psi = E a^- \psi$$

(operators ignore constants like  $E$ ) which is, employing  $a^- H - H a^- = +\hbar\omega a^-$ , we have

$$(+\hbar\omega a^- + H a^-) \psi = E (a^- \psi)$$

$$(+\hbar\omega a^-)\psi + (Ha^-)\psi = E(a^-\psi)$$

$$H(a^-\psi) = E(a^-\psi) - \hbar\omega(a^-\psi)$$

$$H(a^-\psi) = (E - \hbar\omega)(a^-\psi)$$

What does this last statement say? It says that  $a^-\psi$  is *also* an eigenfunction of H if  $\psi$  was. The only thing is that the energy is lower than the energy that was associated with  $\psi$  by  $\hbar\omega$ .

What we have shown is that  $a^-$  is an operator, a ladder operator, which ladders us down!

#### IV. FINDING THE LOWEST EIGENVALUE AND ITS ASSOCIATED EIGENVECTOR

The last part of this wonderful derivation follows. What happens when we continuously “down ladder” from any eigenfunction? Clearly, at some point, we must run out, since the energy can not be lowered indefinitely. Said another way, there must be a lowest eigenfunction, call it  $\psi_0$ , which is destroyed if we ladder down from it.

$$a^-\psi_0 \rightarrow 0$$

Then, we have

$$(p - i\mu\omega x)\psi_0 = 0$$

which means

$$-i\hbar \frac{\partial \psi_0}{\partial x} = i\mu\omega x \psi_0$$

which is immediately integrable. We obtain

$$\psi_0 = C e^{-\frac{\mu\omega}{\hbar} \frac{x^2}{2}} = C e^{-\frac{\mu\sqrt{k\mu}}{\hbar} \frac{x^2}{2}} = C e^{-\frac{\sqrt{k\mu}}{\hbar} \frac{x^2}{2}}$$

If this is the ground state, then laddering up with the up-ladder operator,  $a^+$  should give the first excited state,

$$a^+ C e^{-\frac{\sqrt{k\mu}}{\hbar} \frac{x^2}{2}} = (p + i\mu\omega x) C e^{-\frac{\sqrt{k\mu}}{\hbar} \frac{x^2}{2}}$$

which is

$$-i\hbar \frac{\partial C e^{-\frac{\sqrt{k\mu}}{\hbar} \frac{x^2}{2}}}{\partial x} + i\mu\omega x C e^{-\frac{\sqrt{k\mu}}{\hbar} \frac{x^2}{2}}$$

i.e.,

$$\left(-i\hbar \left(C \left(-\frac{\sqrt{k\mu}}{\hbar} 2x\right)\right) e^{-\frac{\sqrt{k\mu}}{\hbar} \frac{x^2}{2}} + i\mu\omega x C e^{-\frac{\sqrt{k\mu}}{\hbar} \frac{x^2}{2}}\right)$$

which equals

$$\left(i\hbar^2 \frac{\sqrt{k\mu}}{\hbar} 2x + i\mu\sqrt{\frac{k}{\mu}} x\right) C e^{-\frac{\sqrt{k\mu}}{\hbar} \frac{x^2}{2}}$$

or

$$C' x e^{-\frac{\sqrt{k\mu}}{\hbar} \frac{x^2}{2}} \rightarrow \psi_1$$

Continuing, we have

$$a^+ C' x e^{-\frac{\sqrt{k\mu}}{\hbar} \frac{x^2}{2}} \rightarrow \left(-i\hbar \frac{\partial}{\partial x} + i\mu\omega x\right) C' x e^{-\frac{\sqrt{k\mu}}{\hbar} \frac{x^2}{2}}$$

which is, carrying out the differentiation

$$\begin{aligned} &= -i\hbar \frac{\partial C' x e^{-\frac{\sqrt{k\mu}}{\hbar} \frac{x^2}{2}}}{\partial x} + i\mu\omega x C' x e^{-\frac{\sqrt{k\mu}}{\hbar} \frac{x^2}{2}} \\ &= \left(-i\hbar C' + 2i\hbar \frac{\sqrt{k\mu}}{\hbar} C' \frac{x^2}{2} + i\mu\omega C' \frac{x^2}{2}\right) e^{-\frac{\sqrt{k\mu}}{\hbar} \frac{x^2}{2}} \end{aligned}$$

$$iC' \left(-\hbar + \sqrt{k\mu} x^2 + \sqrt{k\mu} \frac{x^2}{2}\right) e^{-\frac{\sqrt{k\mu}}{\hbar} \frac{x^2}{2}}$$

$$iC' \left(-\hbar + \sqrt{k\mu} x^2\right) e^{-\frac{\sqrt{k\mu}}{\hbar} \frac{x^2}{2}}$$

$$iC' \hbar \left(\frac{\sqrt{k\mu}}{\hbar} x^2 - 1\right) e^{-\frac{\sqrt{k\mu}}{\hbar} \frac{x^2}{2}} \rightarrow \psi_2$$

You will recognize, from this result, that life would have been simpler had we used a dimensionless coordinate for the distance, rather than  $x$ . We keep bringing square roots, and  $\hbar$ 's, which could have been eliminated had we used a dimensionless “ $x$ ”.

#### V. MATRIX ELEMENTS USING LADDER OPERATORS

Be that as it may, we see that we can now evaluate common matrix elements in a simple manner now. Consider  $\langle n|x|m \rangle$  which is a transition matrix element associated with selection rules. Since

$$a^+ - a^- = p + i\mu\omega x - (p - i\mu\omega x)$$

We have

$$\frac{a^+ - a^-}{2i\mu\omega} = x$$

so

$$\langle n|x|m \rangle = \langle n | \frac{a^+ - a^-}{2i\mu\omega} | m \rangle$$

and we know what the effect of the ladder operators are on the bra's and ket's involved. once we've expanded this maximally, and used the ladder operator properties all one need do is preserve the  $\langle i||i \rangle$  terms, since the

$\langle i||j \rangle$  terms vanish (orthogonality) when  $i$  is different from  $j$ !

Consider  $\langle n|p_x|m \rangle$  which is a transition matrix element associated with selection rules. Since

$$a^+ + a^- = p + i\mu\omega x + (p - i\mu\omega x)$$

We have

$$\frac{a^+ + a^-}{2} = p_x$$

so

$$\langle n|p_x|m \rangle = \langle n|\frac{a^+ + a^-}{2}|m \rangle$$

Similar schemes can be used for powers of  $x$  or the momentum, or mixed expressions. But the central idea is that when these ladder operators operate on the right hand ket, they generate, aside from constants, other kets, which may or may not be orthogonal to the bra's in the expression. This leads to obvious simplifications.